



3 IN 1 WEED & FEED for SOUTHERN LAWNS

MATA LAS MALEZAS*; PREVIENE LAS MALEZAS* POR 6 MESES; Y FERTILIZA

1 KILLS
WEEDS* INCLUDING DOLLARWEED, DANDELION & CLOVER

2 FEEDS
GREENS & STRENGTHENS YOUR LAWN

3 PREVENTS
BROADLEAF & GRASSY
WEEDS* PLUS CRABGRASS UP TO **6** MONTHS



TREATS UP TO
5,000 Sq Ft
NET WT 12.5 Lbs (5.6 kg)

Active Ingredients:	
Dicamba acid	0.088%
Penoxsulam	0.041%
Indaziflam	0.027%
Other Ingredients	99.844%
Total	100.000%

*LISTED WEEDS

KEEP OUT OF REACH OF CHILDREN
CAUTION See back panel for additional
precautionary statements

35-0-3

FOR USE ON
SOUTHERN TURFS
BERMUDAGRASS, ST AUGUSTINEGRASS,
CENTIPEDEGRASS & ZOYSIA



Tired of killing weeds in your lawn – even after you've put down your traditional weed & feed granules? Just one application of Bayer Advanced® 3-in-1 Weed & Feed for Southern Lawns kills lawn weeds*, and then prevents new ones, including Crabgrass, for up to 6 months. Plus it greens, feeds and strengthens your lawn for a thick turf. It delivers season long weed control that is simple and easy to use and it will not harm your lawn when used as directed. No need for frequent re-application. Plus, when applied in early spring, it prevents Crabgrass and when applied late summer, Annual Bluegrass (Poa annua).

*LISTED WEEDS

**Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.

quick FACTS

- KILLS BROADLEAF WEEDS*** INCLUDING DOLLARWEED, DANDELION, CLOVER, CHICKWEED & HENBIT
- FEEDS & STRENGTHENS THE LAWN.** CONTAINS SLOW RELEASE FERTILIZER FOR LONG LASTING FEED.
- PREVENTS GRASSY AND BROADLEAF WEEDS* SUCH AS CRABGRASS, ANNUAL BLUEGRASS, COMMON DANDELION, GOOSEGRASS & SPURGE UP TO 6 MONTHS**

WHEN TO APPLY

Apply anytime in early spring or fall.

WHERE TO USE

Use only on established lawns of the turf types listed below:

Bermudagrass, Centipedegrass, all varieties of St. Augustinegrass (including Floratam) and Zoysia

DO NOT USE ON ANY OTHER TURF TYPES

including: Bahiagrass, Bentgrass, Ormond variety of Bermudagrass, Buffalograss, Carpetgrass, Dichondra, Fescue, Kikuyugrass, Ryegrass, or Seashore Paspalum lawns.

COVERAGE

This bag treats 5,000 sq. ft. Questions or comments, call toll-free (877) 229-3724 www.BayerAdvanced.com



For the full Spanish translation go to www.BayerAdvanced.com



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEFORE YOU USE • Use this product only on established Southern lawns listed on this label – new lawns must be at least 3 months old if sodded, 16 months old if sprigged & 16 months old if seeded. • Do not apply this product to water-saturated soils. • Apply only to actively growing turf. • Apply in spring after the lawn is actively growing and has been mowed 2 times. Do not apply while turfgrass is just turning green in early spring. • Apply in fall no closer than 6 weeks before the first historical frost date for your area. • Do not apply to Centipedegrass after October 1. • Do not apply this product if the lawn appears unhealthy e.g., due to lack of water (drought, watering restrictions), turf disease (dollarspot, large patch, grey leaf spot, take-all root rot, etc.), parasitic microscopic worms (nematodes), turf insects (chinch bugs, etc.), heavy shade, recent aeration or verticuting, or if the lawn is prone to winter injury. • Do not allow granules to come into contact with flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur. • Apply when the air is calm to avoid drift onto desirable plants.

HOW TO USE • Use rotary or drop type spreader to apply granules evenly over the lawn. To be sure you cover lawn areas uniformly, first treat the border of the lawn then fill in the center. Make parallel passes, walking at a steady pace to ensure even distribution of the granules. • Sweep, brush, or blow any accidentally applied material off driveways, sidewalks etc. back onto the lawn. • Wait 24 hours after applying then activate residual barrier by lightly watering in. • Do not overlap treated areas or make repeat applications within 90 days as over application could damage the lawn. • Do not make more than two applications per site in any 12 month period. • Heavy raking will disturb the weed preventative barrier and reduce the effectiveness of this product.

FOR BEST RESULTS

WHEN TO APPLY • Apply when weeds are small and actively growing - temperatures are between 50-90°F. • May be applied to dry or (for best results) to dew-wetted lawn. • To prevent Crabgrass and Goosegrass apply early spring before the temperatures are regularly in the lower 80°F range. • To prevent Annual Bluegrass (Poa annua) apply August - December. • Lawns may be seeded 12 months, sodded 6 months, and sprigged 12 months after applying this product.

MOWING • If the lawn needs mowing, do it 1 to 2 days before (for best results) or 1 day after applying this product. • Clippings from the first three mowings should be left on the treated area. Do not use the clippings from the treated areas as mulch or compost around flowers, ornamentals, trees, or in vegetable gardens.

WATERING • If lawn is dry, water 1 to 2 days before using this product. • Do not water for at least 24 hours after application or apply if rain is expected within 24 hours.

HOW IT WORKS This product contains a proven blend of weed killers to kill existing weeds* plus a residual herbicide to stop weed growth until weeds* are dead within weeks. Plus, it feeds your lawn for quick greening and long lasting feeding.


Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.



SPREADER SETTINGS

If your spreader is not listed, please call our Consumer Service Hotline toll-free (877) 229-3724 for help.

- Settings are approximate. The condition of your spreader, the speed at which you walk and uneven terrain may affect the rate of application. Calibrate your spreader according to manufacturer's directions.
- Tip: 1,000 sq ft = 20 ft x 50 ft
- Do not use a spreader that applies granules in narrow, concentrated or uneven bands. Be sure the spreader works properly and spreads granules evenly.



SETTING TO DELIVER 2.5lb/1000 sq. ft.

Broadcast Spreader	SETTING
Earthway Ev-N-Spread	10
Scotts	3
Sta-Green	3
Drop Spreader	SETTING
Scotts Accugreen 1000	2.5
Scotts Accugreen 3000	2

KILLS

Beggarweed, Creeping	Dollarweed (Pennywort)	Peppergrass
Beggarweed, Dwarf	Goldenrod	Pepperweed, Virginia
Betony, Florida	Hawkweed	Plantain, Broadleaf
Burdock	Healall	Smartweed
Buttercup, Hairy	Ivy, Ground (Creeping Charlie)	Sorel, Sheep
Buttonweed, Virginia ¹	Knotweed	Speedwell, Common
Clover, Hop	Kyllinga, Green ¹	Spotted Catsear
Clover, Red	Lespedeza, Common	Spurweed
Cocklebur, Common	Mallow, Roundleaf	Thistle
Coffeeweed,	Matchweed	Vetch, Common
(Common Chickory)	Nutsedge, Purple ¹	Wild Violet
Daisy, English (Lawn Daisy)	Nutsedge, Yellow ¹	Woodsorrel, Creeping
Dandelion, False	Parsley-piert	¹ Suppression

PREVENTS

Ageratum, Tropic	Crabgrass, Large	Foxtail, Giant	Mustard, Black	Ryegrass, Perennial	Velvetleaf
Barley, Mouse	(Hairy Crabgrass)	Foxtail, Green	Mustard, Wild	Sandbur	Vetch, Purple
Barryardgrass	Crabgrass, Smooth	Foxtail, Yellow	Nightshade, American Black	Sedge, Annual	Wild Carrot
Bittercress	Cudweed, Linear Leaf	Flaree, Redstem	Nightshade, Hairy	Sesbania, Hemp	Willowherb
Black Nightshade, Hairy	(Purple Cudweed)	Goosegrass	Parthenium	Sida, Prickly	Woodsorrel (Oxalis)
Bluegrass, Annual (Poa annua)	Dandelion, Cat's Ear	Groundsel, Common	Phyllanthus, Longstalked	(Teaweed)	
Brome, Foxtail	Doveweed	Guineagrass	Pigweed, Prostrate	Southern Brassbuttons	
Buckwheat, Wild	Evening Primrose,	Kochia	Pigweed, Redroot	Spodoptera, Annual	
Burweed, Lawn	Common	Kyllinga, Fragrant	Plantain, Paleseed	Speedwell, Corn	
California Burclover	Evening Primrose,	Lovegrass, Tufted	Puncturevine	Spurge, Prostrate	
Chamomile, False	Cutleaf	Mallow, Little	Redmaids	Sunflower, Common	
Cheatgrass	Flatsedge, Rice	Marestail	Rocket, London	Swinecress	
Crabgrass, Henry	Fleabane, Hairy	Mulberry Weed	Ryegrass, Italian	Tassel Flower	

KILLS AND PREVENTS

Burnweed, American	Purslane, Common
Chickweed, Common	Pusley, Florida
Chickweed, Mouseear	Ragweed, Common
Claver, White	Sheepspurge
Dandelion, Common	Spurge, Spotted
Dock, Curly	Thisle, Canada
Eclipta	Woodsorrel, Yellow
Henbit	¹ Suppression
Knotweed, Prostrate	
Kyllinga, Annual ¹	
Lambsquarters	
Medic, Black	
Plantain, Buckhorn	

ENVIRONMENTAL HAZARDS

- This product is toxic to fish, aquatic invertebrates, and plants.
- To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.
- **Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may reach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.
- **Surface Water Advisory:** This pesticide may impact water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

RIESGOS AMBIENTALES • Este producto es tóxico para los peces, organismos invertebrados acuáticos y las plantas. • Para proteger al medio ambiente, no permitir que el pesticida ingrese o corra por los drenajes de tormenta, drenaje de aguas pluviales, canaleras o aguas superficiales. Aplicar este producto en clima calmo cuando no haya pronóstico de lluvia en las próximas 24 horas para asegurar que el viento o la lluvia no soplen o laven el pesticida fuera del área de tratamiento. Barrer todo producto que quede en caminos, aceras o calle, de nuevo hacia el área tratada del césped o jardín ayudará a evitar que el producto ingrese en cuerpos de agua o sistemas de drenaje. • **Asesoramiento sobre aguas subterráneas:** Este pesticida tiene propiedades y características asociadas con químicos detectados en aguas subterráneas. Este químico puede escurrirse a las aguas subterráneas si se usa en suelos donde sea permeable especialmente donde la mesa de agua sea poco profunda. • **Asesoramiento sobre aguas superficiales:** Este pesticida puede impactar sobre la calidad de agua si dreña al agua de lluvia. Este es real para suelos con mal drenaje y suelos con aguas subterráneas poco profundas. Este producto está clasificado como que tiene un alto potencial de alcance al agua de la superficie a través del drenaje durante varios meses o más después de su aplicación.

PRECAUTIONARY STATEMENTS

CAUTION HAZARDS TO HUMANS & DOMESTIC ANIMALS

- Harmful if swallowed or absorbed through skin. • Causes moderate eye irritation.
- Avoid contact with eyes, skin, or clothing. • Remove and wash contaminated clothing before reuse.
- When using this product, wear long-sleeved shirt, long pants, socks, shoes, and rubber gloves.

After using this product:

- Rinse gloves before removing and wash hands. • If product gets on skin, wash thoroughly with soap & water.
- If product gets on clothing, remove clothing as soon as possible and launder clothing separately before reuse.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

DECLARACIONES DE PRECAUCIÓN

PRECAUCIÓN RIESGOS PARA SERES HUMANOS Y ANIMALES DOMÉSTICOS • Es perjudicial si se ingiere o absorbe por la piel. • Causa irritación moderada en los ojos. • Evitar el contacto con los ojos, la piel o la ropa. • Quitar la ropa contaminada y lavarla antes de volver a usar. • Cuando use este producto, use camiseta de mangas largas, pantalones largos, calcetines, calzado y guantes de goma.

Después de usar este producto:

- Enjuagar los guantes antes de quitárselos y lavarse las manos. • Si el producto entra en contacto con la piel, lavarse bien con agua y jabón.
- Si el producto entra en contacto con la ropa, quitarse la ropa tan pronto como sea posible y lavarla por separado antes de volver a utilizar.
- Lavarse bien con agua y jabón después de manipular y antes de comer, beber, consumir goma de mascar, tabaco o usar el baño

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 - 20 minutes.
- Call a poison control center or doctor for further treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for further treatment advice.

PRIMEROS AUXILIOS

EN CASO DE INGESTIÓN • Llamar inmediatamente a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. • Hacer que la persona beba tragos de agua, si puede tragar. • No inducir el vómito salvo que así lo indique el médico o el centro de control de envenenamiento. • No administrar nada por vía oral a una persona inconsciente. **EN CASO DE CONTACTO CON LA PIEL O LA ROPA** • Quitar la ropa contaminada. • Enjuagar la piel inmediatamente con abundante agua durante 15 a 20 minutos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. **EN CASO DE CONTACTO CON LOS OJOS** • Mantener abiertos los ojos y enjuagar de forma lenta y suave con agua durante entre 15 y 20 minutos. • Retirar los lentes de contacto, si los hubiera, después de los primeros 5 minutos y luego continuar enjuagando los ojos. • Llamar a un médico o a un centro de control de envenenamiento para consultas sobre el tratamiento. **NOTA:** Tener la etiqueta del producto a mano al llamar a un centro de control de envenenamiento. Si busca tratamiento en una sala de emergencias o en el consultorio de un médico, lleve esta etiqueta del producto para mostrársela al personal médico. También puede llamar al número gratuito 1-877-227-3763 para solicitar información médica de emergencia. Ingredientes activos: 0.088% de dicamba, (CAS n.°1918-00-9), 0.041% de penoxsulam (CAS n.°219714-96-2) y 0.027% de indaziflam (CAS n.°950782-86-2).

EPA Reg. No. 72155-112
EPA Est. No. indicated by 3rd and 4th digits of the batch number on this package.
(26) = 82757-FL-1 (57) = 071106-GA-003

Bayer Advanced® 3-in-1 Weed & Feed for Southern Lawns 35-0-3

GUARANTEED ANALYSIS
Total Nitrogen (N) 35.0%
35% Urea Nitrogen*
Soluble Potash (K₂O) 3.0%
Sulfur (S) Total 4.6%
3.5% Free Sulfur (S)
1.1% Combined Sulfur (S) **F1514**

Derived From: Urea, Polymer Coated Sulfur Coated Urea, & Potassium Sulfate
*Contains 17.5% slowly available Nitrogen from Polymer Coated Sulfur Coated Urea

Information regarding the contents and levels of metals in this product are available on the Internet at <http://www.aapfco.org/metals.htm>. Check with your county or city government to determine if there are local regulations for fertilizer use.

Produced for: Bayer Advanced, A Business Unit of Bayer CropScience LP, P.O. Box 12014, 2 T. W. Alexander Dr., Research Triangle Park, NC 27709
Bayer, the Bayer Cross and Bayer Advanced® are registered trademarks of Bayer.

STORAGE & DISPOSAL


Do not contaminate water, food, or feed by storage and disposal

PESTICIDE STORAGE Store in original container in a cool, dry place, out of the reach of children, preferably a locked storage cabinet. Protect from freezing.
PESTICIDE DISPOSAL & CONTAINER HANDLING If Empty: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.
If partly filled: Call your local solid waste agency or toll free 1-877-229-3724 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Research and testing have determined that the "Directions for Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions.

ALMACENAMIENTO Y ELIMINACIÓN

No contaminar el agua o los alimentos al almacenar o eliminar
ALMACENAMIENTO DE PESTICIDAS Almacenar en el contenedor original en un lugar fresco y seco, fuera del alcance de los niños, preferentemente en un gabinete bajo llave. Proteger del congelamiento.
ELIMINACIÓN DE PESTICIDAS Y MANIPULACIÓN DE ENVASES Si está vacío: Envase no rellenable. No rellenar ni volver a utilizar este envase. Colocar el envase vacío en la basura o entregarlo para reciclar si fuera posible. Si está parcialmente lleno: Llamar a la agencia local de residuos sólidos o al número gratuito 1-877-229-3724 para obtener indicaciones para la eliminación. Nunca colocar el producto sin utilizar debajo de algún desagüe interno o externo.



160613AW1