

ARMORY™

BENEFICIAL BACILLUS BLEND

CONTAINS NON-PLANT FOOD INGREDIENT

<i>Bacillus subtilis</i>4 X 10 ⁹ cfu/g
<i>Bacillus amyloliquefacaens</i>4 X 10 ⁹ cfu/g
<i>Bacillus licheniformis</i>4 X 10 ⁹ cfu/g
TOTAL BACILLUS1 X 10 ¹⁰ cfu/g

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

Net Weight – 12.5 lbs (200 oz)

Manufactured and guaranteed by:
Blacksmith BioScience
504 Spring Hill Dr #440
Spring Texas 77386
www.blacksmithbio.com

BLACKSMITH
BIOSCIENCE



Intended Uses: Armory is a high concentration of several select beneficial *Bacillus* bacteria. When applied to plants Armory acts as a probiotic and thereby increasing yield, root vigor and top growth.

Compatibility: Armory can be tank mixed with chemical and fertilizers as well as other biologicals.

APPLICATION DIRECTIONS:

For residential use or smaller quantities

Foliar Spray and Soil Drench: Use 0.25 -1 teaspoons of product per gallon of water as general dilution. Apply as a foliar spray or soil drench applications.

Reservoir Application: Use 0.25 - 2 teaspoons per gallon of reservoir. Reapply each time reservoir water is changed.

Greenhouse/Nursery

Soil Drench: Dissolve 1-8 oz per 100 gallons of water and apply to soil as a normal drench (soak pot until growing media is saturated without creating runoff).

Foliar Spray: Dilute 1-8 oz per 100 gallons and apply as a heavy spray allowing for initial run-off. Reapply every 2-12 weeks

Soil Incorporation: Use 1-2 oz per cubic yard of soil or growing media. Blend evenly into growing media via spray, dry blender, or any other appropriate method.

Outdoor Agriculture Production

Soil Application: Apply 1-6 ounces per acre. Apply Armory to the soil as a drench, through irrigation, at transplant or other means to move the product to the plants root system.

Reapply every 1-8 weeks.

Foliar Application: Use 1-6 ounces per acre.

Dissolve in appropriate amount of water and apply as a foliar spray to plants leaves until wet to run off. Reapply every 7-28 days.

In Furrow Application:

Use 0.5 -6 oz of Armory per acre applied over the seed or as a side dressing during furrow application.

Hydroponic Systems: Use 1-6 oz per 100 gallons of tank water. Dissolve into reservoir before, with or after adding nutrients. Reapply Armory each time reservoir water is changed.

Turf: Apply 6-oz, dissolved in water, for the initial application to the soil. Apply as a foliar spray or soil application. Repeat at 2-oz per acre every 4 weeks throughout the growing season.

Composting: Dissolve 1-2 oz of Armory into 100 gallons of water to create solution. Apply solution to compost at a rate of 100 gallons per 40 cubic yards of soil.

Compost Teas: Add 0.25 – 1.00 tsp (1-4 g) of Armory per gallon of final compost tea solution. If creating compost tea, add Armory after the process is complete and just prior to application.

Bed, Field or Row Preparation: Apply 1-4 ounces per acre. Apply Armory to the soil during tilling, bed preparation or at any time before, during or after planting or harvesting. Apply so the product is moved into the area of the soil where decomposition is needed by tilling in or spraying in heavy.

Post Fumigation Application: To rebuild the microbial population in fumigated soils, apply 4-16 oz of Armory per acre through irrigation directly to the area fumigated. Apply product during or just after transplant. Reapply every 2-8 weeks as needed throughout the growing season.

Crops On Which Armory May Be Used:

<p>Legume vegetables Succulent and dried beans and peas such as green, snap, shell, and Lima beans, garbanzo beans, chickpeas, soybeans, dry beans, peas, split peas, lentils, and other legumes, including those grown for seed production</p>
<p>Oilseed crops Canola, castor, coconut, cotton, flax, oil palm, olive, peanut, rapeseed, safflower, sesame, sunflower, soybeans, and other oilseed crops, including those grown for seed production</p>
<p>Cucurbit Vegetables: Cucumbers, melons, gourds, squash, cantaloupe, and other cucurbits</p>
<p>Fruiting Vegetables: Eggplant, sweet peppers, hot peppers, tomatoes, tomatillos and other fruiting vegetables</p>
<p>Herb, Spices & Mints: Sage, rosemary, thyme, mint, peppermint, dill, basil, oregano, medicinal herbs and other herbs and spices</p>
<p>Leafy Vegetables & Cole Crops: Asparagus, broccoli, Brussels sprouts, cabbage, cauliflower, celery, collards, endive, kale, kohlrabi, lettuce, mustard greens, parsley, spinach and other leafy vegetable crops</p>
<p>Legume & Vegetable Crops: Snap and dry beans, lentils, succulent and dry peas, peanuts, soybeans and other legume/vegetable crops</p>
<p>Small Grains: Rye, rice, sorghum, millet and other small grains</p>
<p>Root Tuber & Bulb Crops: Garlic, onions, potatoes, sweet potatoes, carrot, ginger, ginseng, horseradish, turnip, beets, radish and other root-tuber/bulb crops</p>
<p>Berry Crops: Blueberries, blackberry, raspberry, loganberry, huckleberry, gooseberry, elderberry, currant, caneberry and other berry crops</p>
<p>Citrus Orange, grapefruit, lemon, tangerine, tangelo, lime, pummelo and other citrus crops</p>

<p>Corn Field corn, sweet corn, popcorn, silage corn, seed corn, and other corn crops</p>
<p>Grape Wine grapes, table grapes, raisins and other grape crops</p>
<p>Pome Fruit Apple, crab apple, pear, quince, mayhaw and other pome fruit</p>
<p>Stone Fruit Apricot, cherry, nectarine, peach, plum, prune and other stone fruit</p>
<p>Tree Nuts Almond, pistachio, pecan, walnut, filberts and other tree nuts</p>
<p>Nurseries, Greenhouses, Shadehouse and Ornamental Plants Indoor, outdoor, and shade or other cover grown ornamental trees and shrubs, flowering plants, foliage plants, tropical plants, potted plants, potted or cut flowers, bedding plants, forestry seedlings, conifer production for reforestation, fruit trees, vegetables and other crops grown in greenhouses or nurseries</p>
<p>Turf Grass Turf, sod, lawns, golf course (fairways, roughs, greens, tees), grass seed production Including but not limited to: Bluegrass, Bentgrass, Bermudagrass (common & hybrid), Dichondra, Fescue, Orchardgrass, <i>Poa annua</i>, St Augustine grass, Ryegrass, <i>Zoysia</i>, mixtures, and other grasses or ornamental turf</p>
<p>Other Crops Bananas/ plantains, tobacco, hops, watercress, mushrooms, ginseng, coffee, olives, green leafy plants and all crops grown for seed</p>
<p>Tobacco</p>
<p>Strawberry</p>
<p>Tropical Fruits: Avocado, mango, papaya, banana, pineapple and other tropical fruits</p>

STORAGE AND DISPOSAL

Storage: Do not store in extremely high temperatures (above 130F).

Disposal: Completely empty container. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Limited Warranty / Disclaimer: Blacksmith BioScience LLC warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purpose referred to in the directions for use, timing and method of application, weather, watering practices, nature of soil, condition of the crop, incompatibility with other influencing factors in the use of this product are beyond the control of the seller To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein. No other express or implied warranty of the fitness or merchantability is made

Expiration Date: 00-00-000