



WEED & FEED WITH TRIMEC® HERBICIDE

ACTIVE INGREDIENTS:
 2,4-dichlorophenoxyacetic acid0.55%
 (+)-(R)-2-(2-methyl-4-chlorophenoxy)
 propionic acid0.12%
 Dicamba: 3,6-dichloro-o-anisic acid0.05%
INERT INGREDIENTS:99.28%
TOTAL100.00%

THIS PRODUCT CONTAINS:
 11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
 2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
 1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
 Isomer Specific by AOAC Methods
 TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

EPA Reg. No. 2217-643-32802
 EPA Est. No. 59144-GA-1(A), 32802-MO-1(B),
 37351-CO-1 (C)

Manufactured for: Howard Johnson's Enterprises, Inc.
 9675 S 60th Street, Franklin, WI 53132

Net Weight 50 LBS. (22.68 kg)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STOP!
**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

**PRECAUTIONARY STATEMENTS
Hazards to Humans &
Domestic Animals**

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material.

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

GUARANTEED ANALYSIS

25-00-08
 Total Nitrogen (N) 25%
 25.00% Urea Nitrogen*
 Soluble Potash (K2O) 8%

Derived From: MURIATE OF POTASH,
POLYMER COATED UREA, UREA

* 5.00% Slowly Available Nitrogen from
Polymer Coated Urea

Chlorine (Cl), not more than 6.00%

F315

Net Weight Bulk Density Item Number
 50 lbs (22.70 kg) 52 FGB61861

SAP 100516199

FIRST AID

If on skin or on clothing:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth, if possible. Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information in case of emergency, call toll free 1-877-800-5556.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Do not apply this product to dichondra or carpetgrass. Do not apply this product to 'Floritam' St. Augustinegrass in Florida.
- Do not use this product where desirable clovers are present. Do not exceed specified dosages for any area. Apply when the air is calm. Do not apply to newly seeded lawns. During application, close spreader openings when turning to prevent over-application.
- Do not use this product on or near desirable plants, and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Pennington Seed Weed & Feed with Trimec® Herbicide controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns. In addition, a XX-XX-X fertilizer provides plant nutrients or a supply of plant food. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days. A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing (*Number of pounds) of nitrogen per 1,000 square feet. Please consult your local extension office for questions regarding a regular fertilizer program suitable for your geographic region. The maximum broadcast application rate is 5.0 pounds of product per 1,000 sq.ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 10.0 pounds of product per 1,000 sq.ft.

- Mow lawn to normal height 1 to 2 days before application.
- Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapf-co.org/metals.htm>.

SUGGESTED SPREADER SETTINGS

Spreader setting to apply at the listed rate		
3.33 lbs per 1000 sq ft	Cone setting	
Accu-Pro 2000	6	L 1/2
SR 2000	6	L 1/2
Lesco 021820	0.25 closed	D 1/2
Spyker 76-22		4.3
PSB bigfoot	A.5	J 1/2
Lely HR	11-B	5.25
Vicon 203		15

As a guide, set spreader at the setting indicated by the above chart before applying. Remember the settings provided are only a guide and depend on the age, condition and speed of operation of your spreader. Check spreader calibration often and make adjustments as needed. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before finishing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete. Calibration Instructions: To provide proper distribution, calibrate spreader before application.

The above spreader settings are approximate. 50 lb. bag: Calibrate your spreader on 300 sq.ft. and adjust to apply 1.0 lb. of product per 300 sq.ft

For 50 lb bag: Apply 3.33 lbs of product per 1,000 sq ft of lawn area. Check your spreader owner's manual for calibration settings and set spreader to apply at the rate of 1 pound of product per approximately 300 sq ft. A uniform distribution of product on the treated area is essential. One bag covers 15,000 sq ft.

Choose One Of These Spreader Patterns

- Single Pass** - Operate the equipment at intervals corresponding to the width or swath of the spreader. For drop spreaders, overlap the wheel tracks to ensure complete coverage. Close the spreader openings when turning to change directions.
- Double Pass** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. Close the spreader openings when turning to change directions.
- Cross Hatching** - Reduce the application rate by one-half of the recommended rate, calibrate the spreader, and adjust the settings for the desired application rate. Operate the spreader over the lawn two (2) times. For the second trip, operate the spreader perpendicular to the wheel tracks of the spreader pattern of the first trip. Close the spreader opening when turning to change directions.

Weed & Feed effectively controls the following weeds when used as directed.

WEEDS			
Bedstraw	Ground ivy	Peppergrass	Spurge
Black medic	Healall	Pigweed	Thistle
Buckhorn	Henbit	Plantain	Wild carrot
Burdock	Knotweed	Poison ivy	Wild garlic
Chicory	Lambsquarters	Poison oak	Wild lettuce
Chickweed	Lespedeza	Purslane	Wild onion
Clover	Mallow	Ragweed	Yarrow
Dandelion	Morningglory	Sheep sorrel	
Dock	Oxalis	Speedwell	

**LIMITED WARRANTY AND DISCLAIMER
FOR USE ONLY AS DIRECTED.**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.